

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate only, other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (07804-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (LEAVE BLANK)		2. REPORT DATE  23 April 1999		3. REPORT TYPE AND DATES COVERED  Abstract
4. TITLE AND SUBTITLE  The Effect of Aircrew Age on +Gz Tolerance as Measured in a Human-Use Centrifuge			5. FUNDING NUMBERS	
6. AUTHOR(S)  Estrella Forster      Barry Shender				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Naval Air Warfare Center Aircraft Division 22347 Cedar Point Road, Unit #6 Patuxent River, Maryland 20670-1161			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  Naval Air Systems Command 47123 Buse Road, Unit IPT Patuxent River, Maryland 20670-1547			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES			19991004 318	
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.				
12b. DISTRIBUTION CODE				
13. ABSTRACT (Maximum 200 words)  A data repository was established in 1988 to compile information on 1,120 aircrew (74% pilots) who underwent acceleration (+Gz) tolerance training at NAWCAD Patuxent River human-use centrifuge. 51% of the aircrew flew in high performance aircraft (F4, F14, F15, F16, and F18). The trainees were U.S. Navy/Marine Corps (70%) and Air National Guard (30%) aircrew. Balanced data from 817 healthy male trainees were examined. Mean age (+S.D.) was $31.4 \pm 6.8$ years (20 to 59). Relaxed tolerance was $4.91 \pm 0.93$ +Gz and was shown to be independent of age ( $R^2 = 0.005$ ). Straining tolerance was $7.17 \pm 1.27$ +Gz. Age did not have an effect on straining +Gz tolerance ( $R^2 = 0.017$ ). The protection afforded by the AGSM was $2.72 \pm 0.84$ +Gz and was not affected by trainee age ( $R^2 = 0.007$ ). Age did not demonstrate to have an effect on G-LOC incidence. Exposures where cardiovascular data was analyzed ( $n=19$ ) ranges from 5 to 9 +Gz ( $5.5 \pm 1.3$ ). The change described by MHR-RHR was $57 \pm 21$ bpm. The change described by MHR-RCVHR was $62 \pm 27$ bpm. Multiple regress demonstrated that age and the +Gz level at which the MHR occurred (GMHR) explained 55% of the variability in MHR-RCVHR ( $R^2$ age = 0.18, $pT_{\beta_1} = 0.01$ ; $R^2$ GMHR = 0.37, $pT_{\beta_2} = 0.002$ ). The model was described by $MHR-RCVHR = 19.03 - 1.40 \cdot \text{age} + 13.08 \cdot \text{GMHR}$ ( $F = 9.87$ , $p = 0.001$ ). No statistically significant relationship was found based on change in MHR-RHR. The relative long duration GOR exposures are typically used to determine cardiovascular +Gz tolerance in human-use centrifuge studies. Based on the variables examined in this retrospective study, there does not seem to be a significant effect on age on +Gz tolerance.				
14. SUBJECT TERMS  +Gz tolerance training			15. NUMBER OF PAGES  2	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT  Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE  Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT  Unclassified	20. LIMITATION OF ABSTRACT  UL	

HFM SYMPOSIUM  
On  
"Operational Issues of Aging Crew Members"

THE EFFECT OF AIRCREW AGE ON +Gz TOLERANCE  
AS MEASURED IN A HUMAN-USE CENTRIFUGE

Estrella M. Forster, Ph.D.  
Barry S. Shender, Ph.D.\*

Naval Air Warfare Center Aircraft Division  
Crew Systems Technology Department  
South Engineering Center Bldg. 2187  
Unit 5 Code 4.6.4.1  
48110 Shaw Road  
Patuxent River, MD 20670-1906

Voice: 301-342-9278  
Fax: 301-342-9305  
Email: ForsterEM@navair.navy.mil

\*Voice: 301-342-8881  
Fax: 301-342-8876  
Email: ShenderBS@navair.navy.mil

**ABSTRACT ONLY**

**CLEARED FOR  
OPEN PUBLICATION**

*23 Apr 99*

**PUBLIC AFFAIRS OFFICE  
NAVAL AIR SYSTEMS COMMAND**

*H. Howard*

**INTRODUCTION.** A data repository was established in 1988 to compile information on 1,120 aircrew (74% pilots) who underwent acceleration (+Gz) tolerance training at the Naval Air Warfare Center Aircraft Division human-use centrifuge. 51% of the aircrew flew in high performance aircraft (F4, F14, F15, F16, and F18). The trainees were US Navy/Marine Corps (70%) and Air National Guard (30%) aircrew. **METHOD.** The database was examined to determine the effect of age on +Gz tolerance as measured during gradual onset rate (GOR) exposures. GOR were the first in a series of +Gz exposures during a single training day. GOR commenced at a resting level of 1.0 +Gz. +Gz then increased at 0.1 G/s until the trainee experienced 60° Peripheral Light Loss (PLL1) as defined by the inability to see an array of LEDs placed in an arc describing 15° increments (150° total) 30 cm in front of the trainee at eye level. Once PLL1 was reached, trainees performed Anti-G Straining Maneuvers (AGSM) until 60° PLL was reached again (PLL2). Trainees then terminated the +Gz exposure by pressing a button located on a control stick. The limit of the exposures was 9 +Gz. The variables selected for analysis were: trainee relaxed +Gz tolerance (+Gz load at PLL1); trainee straining tolerance (+Gz load at PLL2); and the protection afforded by the AGSM (PLL2-PLL1). Incidence of G-induced Loss of Consciousness (G-LOC) with respect to age was also examined. Resting (RHR), maximum (MHR), and recovery heart rate (RCVHR) available from 19 subjects was also examined to determine the effect of age on baroreceptor response to +Gz and recovery to resting levels.

**RESULTS.** Balanced data from 817 healthy male trainees were examined. Mean age ( $\pm$  S.D.) was  $31.4 \pm 6.8$  years (20 to 59). Relaxed tolerance was  $4.91 \pm 0.93$  +Gz and was shown to be independent of age ( $R^2 = 0.005$ ). Straining tolerance was  $7.17 \pm 1.27$  +Gz. Age did not have an effect on straining +Gz tolerance ( $R^2 = 0.017$ ). The protection afforded by the AGSM was  $2.72 \pm 0.84$  +Gz and was not affected by trainee age ( $R^2 = 0.007$ ). Age did not demonstrate to have an effect on G-LOC incidence. Exposures where cardiovascular data was analyzed ( $n = 19$ ) ranged from 5 to 9 +Gz ( $5.5 \pm 1.3$ ). The change described by MHR-RHR was  $57 \pm 21$  bpm. The change described by MHR-RCVHR was  $62 \pm 27$  bpm. Multiple regression demonstrated that age and the +Gz level at which the MHR occurred (GMHR) explained 55% percent of the variability in MHR-RCVHR ( $R^2$  age = 0.18,  $p_{T_{p1}} = 0.01$ ;  $R^2$  GMHR = 0.37,  $p_{T_{p2}} = 0.002$ ). The model was described by  $MHR-RCVHR = 19.03 - 1.40 * age + 13.08 * GMHR$  ( $F = 9.87$ ,  $p = 0.001$ ). No statistically significant relationship was found based on change in MHR-RHR. **CONCLUSION.** The relatively long duration GOR exposures are typically used to determine cardiovascular +Gz tolerance in human-use centrifuge studies. Based on the variables examined in this retrospective study, there does not seem to be a significant effect of age on +Gz tolerance.